

5. Properties of Abstract Voronoi Diagrams


5.1 Euclidean Voronoi Diagrams

Voronoi Diagram: Given a set S of n point sites in the plane, the Voronoi diagram $V(S)$ of S is a planar subdivision such that

- Each site $p \in S$ is assigned a Voronoi region denoted by $\text{VR}(p, S)$
- All points in $\text{VR}(p, S)$ share the same nearest site p in S


Voronoi Edge: The common boundary between two adjacent Voronoi regions, $\text{VR}(p, S)$ and $\text{VR}(q, S)$, i.e., $\text{VR}(p, S) \cap \text{VR}(q, S)$, is called a *Voronoi edge*.

Voronoi Vertex: The common vertex among more than two Voronoi regions is called a *Voronoi vertex*.




The Euclidean Voronoi diagram can be computed in $O(n \log n)$ time


- Line Segment Voronoi Diagram



- Circle Voronoi Diagram



- Voronoi Diagram in the L_1 metric



5.2 Bisecting Systems

- For two sites $p, q \in S$, the bisector $\mathbf{J}(p, q)$ between p and q is defined as $\{x \in R^2 \mid d(x, p) = d(x, q)\}$
- $J(p, q)$ partitions the plane into two half-planes
 - $D(p, q) = \{x \in R^2 \mid d(x, p) < d(x, q)\}$
 - $D(q, p) = \{x \in R^2 \mid d(x, q) < d(x, p)\}$
- $\text{VR}(p, S) = \bigcap_{q \in S \setminus \{p\}} D(p, q)$
- $V(p, S) = R^2 \setminus \bigcup_{p \in S} \text{VR}(p, S)$
 - consists of Voronoi edges.

5.3 Abstract Voronoi Diagrams


A unifying approach to computing Voronoi diagrams among different geometric sites under different distance measures.

A bisecting system $\mathcal{J} = \{J(p, q) \mid p, q \in S\}$ for a set S of sites (indices)


A bisecting system \mathcal{J} is **admissible** if \mathcal{J} satisfies the following axioms

- (A1) Each bisecting curve in \mathcal{J} is homeomorphic to a line (not closed)
- (A2) For each non-empty subset S' of S and for each $p \in S'$, $\text{VR}(p, S')$ is path-connected.
- (A3) For each non-empty subset S' , $R^2 = \bigcup_{p \in S'} \overline{\text{VR}(p, S')}$
- (A4) Any two curves in \mathcal{J} have only finitely many intersection points, and these intersections are transversal.

- (A1) can be written as "Each curve in \mathcal{J} is unbounded. After stereographic projection to the sphere, it can be completed to a closed Jordan curve through the north pole."
- (A4) can be removed through several complicated proofs.




Not Traversal




Traversal

Not Admissible




Non-Man Land



Disconnected

Three possibilities of an admissible system for three sites



Abstract Voronoi Diagrams

- A category of Voronoi diagrams
 - points in any convex distance function
 - Karlsruhe metric
 - Line segments and convex polygons of constant size

5.3 Basic Properties

Lemma 1

Let (S, \mathcal{J}) be a bisecting curve system. The the following assertions are equivalent.

1. If $p, q,$ and r are pairwise different sites in S , then $D(p, q) \cap D(q, r) \subseteq D(p, r)$ (Transitivity)
2. For each nonempty subset $S' \subseteq S$, $R^2 = \bigcup_{p \in S'} \overline{\text{VR}(p, S')}$

Proof:

(2) \rightarrow (1)

- Let z be a point in $D(p, q) \cap D(q, r)$.
- By (2), there must be a site $t \in S' = \{p, q, r\}$ such that $z \in \text{VR}(t, S')$.
- If $t = p$, $z \in \text{VR}(p, S') \subseteq D(p, r)$; otherwise
 - $z \in \text{VR}(q, S') \subseteq D(q, p)$, contradicting $z \in D(p, q)$
 - $z \in \text{VR}(r, S') \subseteq D(r, q)$, contradicting $z \in D(q, r)$

(1) \rightarrow (2)

- By induction on $|S'|$.
- If $|S'| = 2$, the assertion is immediate.
- The case where $|S'| = 3$ follows directly from (1)
- Let z be a point in the plane. By induction hypothesis, to each $p \in S'$, there exists a site $c(p) \neq p$ such that $z \in \text{VR}(c(p), S' \setminus \{p\})$

case 1: There exists $v \neq w$ such that $c(v) = c(w)$. Then

$$\begin{aligned} z &\in \text{VR}(c(v), S' \setminus \{v\}) \cap \text{VR}(c(v), S' \setminus \{w\}) \\ &\subset \text{VR}(c(v), S' \setminus \{v\}) \cap D(c(v), v) = \text{VR}(c(v), S') \end{aligned}$$

case 2 The mapping c is injective. Let p, v, w be such that $|\{p, c(p), v, w\}| = 4$. Since $c(v) \neq c(w)$, one of them is different from p . We assume $c(v)$ is different from p . Since $c(v) \neq c(p)$ we obtain the contradiction:

$$\begin{aligned} z &\in \text{VR}(c(p), S' \setminus \{p\}) \subseteq D(c(p), c(v)) \\ z &\in \text{VR}(c(v), S' \setminus \{v\}) \subseteq D(c(v), c(p)) \end{aligned}$$

Theorem



A bisecting curve system (S, \mathcal{J}) is *admissible* if and only if the following conditions are fulfilled.

1. $D(p, q) \cap D(q, r) \subseteq D(p, r)$ holds for any three sites p, q, r , in S
2. Any two curve $J(p, q)$ and $J(p, r)$ cross at most twice and do not constitute a clockwise cycle in the plane

proof

→

- By Lemma 1, concentrate on the connectedness of Voronoi regions.
- Consider an infinitely large bounded curve Γ which contains all intersections among curves in \mathcal{J}
- For any $p, q, r \in S$, $V(\{p, q, r\})$ encircled by Γ is a planar graph with exactly 4 faces each of whose vertices is of degree at least 3.
- By the Euler Formula, the planar graph has at most 4 vertices
- Since at least two edges of the original diagram tend to infinity, two vertices must be situated in Γ .
- $J(p, q)$ and $J(p, r)$ cross at most twice since each intersection between them is a Voronoi vertex by definition.
- A simple case analysis shows no clockwise cycle arising from $J(p, q)$ and $J(p, r)$



←

- The case analysis shows that for any 3-element subset S' of S , all Voronoi regions in $V(S')$ is connected.
- We prove by induction on m : If $R = \text{VR}(p, \{p, q_1, q_2, \dots, q_m\})$ is connected, then $R \cap D(p, q_{m+1}) = \text{VR}(p, \{p, q_1, q_2, \dots, q_{m+1}\})$ is connected.
- Let $J(p, q)$ be oriented such that $D(p, q)$ is on its left side.
- Assume the contrary that $R \cap D(p, q_{m+1})$ were not connected.
- If $R \cap D(p, q_{m+1})$ is bounded, let C be ∂R and $J(p, q_{m+1})$ would form a clockwise cycle.
 - For $\exists i \leq m$, $J(p, q_i)$ and $J(p, q_{m+1})$ form a clockwise cycle.
 - There exists a contradiction
- Otherwise, we intersect R with the inner domain of Γ , and C' be its contour.
 - The same reasoning applies to C' and $J(p, q_{m+1})$

