

Exercise 23: Shatter Function Lemma (4 Points)

1. Show the correctness of

$$\binom{m-1}{i} + \binom{m-1}{i-1} = \binom{m}{i}.$$

2. Show that the bound (ii) in the Shatter Function Lemma is tight! Construct a set system \mathcal{F} for all d and m such that $VCDim(\mathcal{F}) = d$ and $\pi_{\mathcal{F}}(m) = \Phi_d(m)$, where $\Phi_d(m) = \binom{m}{0} + \binom{m}{1} + \dots + \binom{m}{d}$ holds.
3. Clarify the proof detail on page 111 of the manuscript:

$$\left(1 - \frac{d}{m}\right)^{d-m}$$

is increasing in m !




Figure 1: The points p and q_1 are L_1 -visible whereas p and q_2 are not L_1 -visible because the L_1 -visibility is blocked by the horizontal L_1 -cut of the locally Y -minimal vertex v_2 .

Exercise 24: VC Dimension L_1 -visibility (4 Points)

Consider the following notion of L_1 -visibility inside a simple polygon P : Two points p and q of inside P are L_1 -visible to each other inside P , iff there is an L_1 -path inside P from p to q that is *monotone* in X - and Y -direction, see the Figure for some examples.

Try to find an example in order two show that the VC-Dimension of points in simple polygons is 3 (or even 4) w.r.t. L_1 -visibility polygons of P !